

Monitoring Earth's global energy balance from space

– requirements –

Thorsten Mauritsen, Steven Dewitte, Thomas Hocking, Linda Megner, Luca Schifano





Royal Observatory
of Belgium



Stockholms
universitet

Climate out of balance

- Energy is accumulating because of increasing greenhouse gases
- This causes:
 - Rising temperatures
 - Shifting climate zones
 - Droughts and heat waves
 - Flooding
 - Increasing sea levels
- Yet, imbalance is only 1 out of 340 Wm^{-2} , or **0.3 percent**



Royal Observatory
of Belgium





Stockholms
universitet

Societal benefits from long term monitoring

Governments of the world have agreed to limit global warming below 2 degrees

- Requires declining imbalance:
 - follow up on international agreement
- Prepare for worst-case scenarios:
 - aerosol forcing larger than expected: rapid warming ahead!
 - global tipping-point, extremely unlikely but catastrophic impact
 - geo-engineering may be needed in future

Simulations of past and future scenarios:



Royal Observatory
of Belgium



Stockholms
universitet

Requirements

To be useful, a system must have drifts and errors that are smaller than the signal we want to measure, preferably much smaller

Drift	$\ll 0.2 \text{ Wm}^{-2} / \text{decade}$
Systematic error	$< 1.0 \text{ Wm}^{-2}$
Random error	$< 1.0 \text{ Wm}^{-2}$



Royal Observatory
of Belgium



Stockholms
universitet

Making the problem simpler

- Sacrifice resolution for accuracy
- Focus on global annual mean imbalance
- Maintainable and stable over decades





Royal Observatory
of Belgium



Stockholms
universitet

Making the problem simpler

- Sacrifice resolution for accuracy
- Focus on global annual mean imbalance
- Maintainable and stable over decades



Schifano et al. (2020)



Royal Observatory
of Belgium




Stockholms
universitet


Past, current and planned missions use scanners

All scanning radiometers depend on angular dependency models (ADMs) to correct for narrow field of view

- ERBE
- CERES
- Libera
- CLARREO-Pathfinder
- ESAs TRUTHS, FORUM, IASI-NG



$$F(\theta_s) = \frac{\pi I(\theta_s, \theta_v, \phi)}{R(\theta_s, \theta_v, \phi)}$$



From Gristey et al. (2021)



Royal Observatory
of Belgium




Stockholms
universitet

Anisotropic effects

Wide field of view radiometer measures actual flux at satellite position



However, systematic biases in viewing- and solar zenith angles may introduce biases:

- Gristey et al. (2017) explored effects of anisotropic using TRMM angular dependence model (ADM)
- Found difference when introducing ADM of 1.6 Wm^{-2} compared with isotropic case
- But only 0.1 Wm^{-2} between true and randomised ADMs



The Earth Climate Observatory (ECO) mission

- Consist of polar orbiting satellites
- Observe incoming and outgoing radiation with identical instruments
- Rotate to cancel systematic calibration errors
- Spare instruments to monitor slow drift
- Wide angle cameras (solar/terrestrial) for separation, scene identification, ADM development, mapping
- Constellation to improve sampling, possibly combining precessing and sun-synchronous orbits




Royal Observatory
of Belgium




Stockholms
universitet

The Earth Climate Observatory (ECO) mission




1- 2 satellites in precessing orbits:

- Good sampling of diurnal cycle on annual time scale
- Only annual means
- Single point of failure
- Mapping difficult




3-4 sun-synchronous + 1-2 precessing:

- Excellent sampling of diurnal cycle on annual time scale
- Cloud feedback monitoring
- Insensitive to failure



2 sun-synchronous + 1-2 precessing:

- Good sampling of diurnal cycle on annual time scale
- Monthly means + mapping possible
- Intercalibration
- Sensitive to failure



8, or more, sun-synchronous:

- Excellent sampling of diurnal cycle on daily time scale
- Footprints overlap, also in tropics
- Excellent mapping
- Robust to failure



Royal Observatory
of Belgium




Stockholms
universitet

Summary of errors

ERBE non-scanner (from Wong et al. 2018):

Sources of uncertainty	Longwave	Shortwave
Instrument stability (drift)	± 0.5	± 0.1
Instrument absolute accuracy	± 2.5	± 2.5
Intercalibration (footprint mismatch)	± 1.2	± 1.0
Non-scanner inversion (mapping to TOA level)	$< \pm 1.0$	$< \pm 1.0$
Satellite altitude correction	0.0	0.0
Twilight data	n/a	$> \pm 0.2$



~ 3.0 Wm⁻² per channel

From SRL-2 draft report



Royal Observatory
of Belgium



Stockholms
universitet


Summary of errors

ERBE non-scanner (from Wong et al. 2018):

Sources of uncertainty	Longwave	Shortwave
Instrument stability (drift)	± 0.5	± 0.1
Instrument absolute accuracy	± 2.5	± 2.5
Intercalibration (footprint mismatch)	± 1.2	± 1.0
Non-scanner inversion (mapping to TOA level)	$< \pm 1.0$	$< \pm 1.0$
Satellite altitude correction	0.0	0.0
Twilight data	n/a	$> \pm 0.2$

Preliminary ECO mission estimates:

Sources of uncertainty	EEI
Instrument stability (drift per decade)	$\ll \pm 0.1$
Instrument absolute accuracy on difference (preliminary)	$< \pm 0.5$
Intercalibration (footprint mismatch)	n/a
Non-scanner inversion (mapping and reference level)	n/a
Satellite altitude correction	0.0
Twilight data	n/a
Polar regions	?
Diurnal cycle sampling (preliminary estimate)	$\ll \pm 0.3$
Anisotropic correction (literature estimate)	$< \pm 0.1$





~ 3.0 Wm⁻² per channel

Long term monitoring strategy

ECO mission nominal life time is relatively short (5-10 years), but the idea is that it can develop into a long term monitoring mission:

- Instruments are fairly simple, and most of the cost is in development
- Spare instruments can serve to:
 - Evaluate issues on ground
 - Quickly launch satellites in case of failure
 - Piggyback on other missions, synergies
 - Help improve future missions
 - Be shared with other space agencies
- Challenge: long term monitoring is not so cool

GLOBAL AVERAGE SURFACE TEMPERATURE



Royal Observatory
of Belgium



Stockholms
universitet

Project status



We are working towards answering an ESA Earth Explorer call next year

Phase F	9	Science Impact Quantified
Phase E2	8	Science Validated and Matured
Phase E1	7	Science Demonstrated
Phase B, C, D	6	Mission Concept Validated
Phase A/B	5	Mission Performance Assessed
Phase 0	4	Feasibility Shown
Development Phase	3	Requirements Drafted
Development Phase	2	Scientific Idea Consolidated
Development Phase	1	Initial Scientific Idea Formulated



Summary ECO mission

- We aim to provide accurate and robust long term monitoring of Earth's **global mean radiation imbalance**
- Concept based on wide field of view radiometers using differential technique plus two cameras (Steven's talk)
- Use of multiple identical instruments to reduce errors from calibration and drift
- Complementary, to more resolution-focussed "big" missions (spatial, temporal, spectral)
- [long list of things it cannot do]
- A 'gap' filler?



Extra slides



Royal Observatory
of Belgium



Stockholms
universitet

Anisotropic effects


Wide field of view radiometer measures actual flux at satellite position

However, systematic biases in viewing- and solar zenith angles may introduce biases:

- Gristey et al. (2017) simulated a 32 satellite constellation to produce hourly maps
- Explored effects of anisotropic using TRMM angular dependence model (ADM)
- Found difference when introducing ADM of 1.6 Wm^{-2} compared with isotropic case
- But only 0.1 Wm^{-2} between true and randomised ADMs

$$\pi^{-1} \int_0^{2\pi} d\phi \int_0^{\pi/2} d\theta R(\theta_0, \theta, \phi) \cos \theta \sin \theta = 1$$

Mean: 0.98



Royal Observatory
of Belgium




Stockholms
universitet

Anisotropic effects

$$\pi^{-1} \int_0^{2\pi} d\phi \int_0^{\pi/2} d\theta R(\theta_0, \theta, \phi) \cos \theta \sin \theta = 1$$

- We currently think the effect is small ($\sim 0.1 \text{ Wm}^{-2}$)
- In this case we can apply a climatological ADM derived from cameras to correct for this small error
- In the unlikely event the effect is large, a correction using scene dependent ADMs may be needed

Mean: 0.98



Royal Observatory
of Belgium



Stockholms
universitet